

# Simulating full QCD at nonzero density using the Complex Langevin Equation

Dénes Sexty

Heidelberg University

Lattice 2013, 31.07.2013.

Collaborators: Gert Aarts (Swansea), Lorenzo Bongiovanni (Swansea)  
Erhard Seiler (MPI München), Ion Stamatescu (Heidelberg)

1. Introduction
2. Gauge symmetry and gauge cooling
3. HQCD with gauge cooling
4. Extension to Full QCD

Seiler, Sexty, Stamatescu PLB (2012)

Aarts, Bongiovanni, Seiler, Sexty, Stamatescu EPJA (2013)

Sexty, arXiv:1307.7748

---

---

# Non-zero chemical potential

Euclidean gauge theory with fermions:  $Z = \int dU \exp(-S_E) \det(M)$

For nonzero chemical potential, the fermion determinant is complex

Sign problem  $\longrightarrow$  Naïve Monte-Carlo breaks down

Methods going around the problem work for  $\mu = \mu_B/3 < T$

(Multi parameter) reweighting

Barbour et. al. '97; Fodor, Katz '01

Analytic continuation of results obtained at imaginary  $\mu$

Lombardo '00; de Forcrand, Philipsen '02; D'Elia Sanfilippo '09; Cea et. al. '08-

Taylor expansion in  $(\mu/T)^2$

de Forcrand et al. '99; Hart, Laine, Philipsen '00; Gavai and Gupta '08;  
de Forcrand, Philipsen '08

Stochastic quantisation

Aarts and Stamatescu '08

Bose Gas, Spin model, etc. Aarts '08, Aarts, James '10 Aarts, James '11

QCD with heavy quarks: Seiler, Sexty, Stamatescu '12

# Stochastic Quantization Parisi, Wu (1981)

Weighted, normalized average: 
$$\langle O \rangle = \frac{\int e^{-S(x)} O(x) dx}{\int e^{-S(x)} dx}$$

Stochastic process for  $x$ : 
$$\frac{dx}{d\tau} = -\frac{\partial S}{\partial x} + \eta(\tau)$$

Gaussian noise  $\langle \eta(\tau) \rangle = 0$   $\langle \eta(\tau) \eta(\tau') \rangle = 2\delta(\tau - \tau')$

Averages are calculated along the trajectories:

$$\langle O \rangle = \frac{1}{T} \int_0^T O(x(\tau)) d\tau$$

Fokker-Planck equation for the probability distribution of  $P(x)$ :

$$\frac{\partial P}{\partial \tau} = \frac{\partial}{\partial x} \left( \frac{\partial P}{\partial x} + P \frac{\partial S}{\partial x} \right) = -H_{FP} P$$

Real action  $\rightarrow$  positive eigenvalues

for real action the Langevin method is convergent

# Langevin method with complex action

Klauder '83, Parisi '83, Hueffel, Rumpf '83,  
Okano, Schuelke, Zeng '91, ...  
applied to nonequilibrium: Berges, Stamatescu '05, ...

$$\frac{dx}{d\tau} = -\frac{\partial S}{\partial x} + \eta(\tau)$$

The field is complexified

real scalar  $\longrightarrow$  complex scalar

link variables: SU(N)  $\longrightarrow$  SL(N,C)  
compact  $\longrightarrow$  non-compact

$$\det(U) = 1, \quad U^\dagger \neq U^{-1}$$

Distance from SU(N)

$$\sum_{ij} |(U U^\dagger - 1)_{ij}|^2$$

Unitarity Norms:

$$\text{Tr}(U U^\dagger) \geq N$$

$$\text{Tr}(U U^\dagger) + \text{Tr}(U^{-1} (U^{-1})^\dagger) \geq 2N$$

For SU(2):  $(\text{Im Tr } U)^2$

Analytic observables

$$\frac{1}{Z} \int P_{\text{comp}}(x) O(x) dx = \frac{1}{Z} \int P_{\text{real}}(x, y) O(x + iy) dx dy$$

# Gaugefixing in SU(2) one plaquette model

Berges, Sexty '08

SU(2) one plaquette model:  $S = i\beta \text{Tr} U \quad U \in \text{SU}(2)$

Langevin updating  $U' = \exp(i\lambda_a(\epsilon i D_a S[U] + \sqrt{\epsilon} \eta_a)) U$

exact averages by numerical integration:  $\langle f(U) \rangle = \frac{1}{Z} \int_0^{2\pi} d\phi \int d\Omega \sin^2 \frac{\phi}{2} e^{i\beta \cos \frac{\phi}{2}} f(U(\phi, \hat{n}))$

“gauge” symmetry:  $U \rightarrow W U W^{-1}$       complexified theory:  $U, W \in \text{SL}(2, \mathbb{C})$


After each Langevin timestep: fix gauge condition

$$U = a \mathbf{1} + i \sqrt{1 - a^2} \sigma_3$$


$$b_i = (0, 0, \sqrt{1 - a^2})$$

# SU(2) one-plaquette model

Distributions of  $\text{Tr}(U)$  on the complex plane



Without gaugefixing



With gaugefixing

Exact result from integration:  $\langle \text{Tr } U \rangle = i 0.2611$

From simulation:

$$(-0.02 \pm 0.02) + i(-0.01 \pm 0.02)$$

$$(-0.004 \pm 0.006) + i(0.260 \pm 0.001)$$

**With gauge fixing, all averages are correctly reproduced**

# Gauge cooling

complexified distribution with slow decay  $\longrightarrow$  convergence wrong results

Minimize unitarity norm:  $\sum_i \text{Tr}(U_i U_i^+)$

Using gauge transformations in  $SL(N, \mathbb{C})$

$$U_\mu(x) \rightarrow V(x) U_\mu(x) V^{-1}(x + a_\mu) \quad V(x) = \exp(i \lambda_a v_a(x))$$

$v_a(x)$  is imaginary (for real  $v_a(x)$ , unitarity norm is not changed)

Gradient of the unitarity norm gives steepest descent

$$G_a(x) = 2 \text{Tr} [\lambda_a (U_\mu(x) U_\mu^+(x) - U_\mu^+(x - a_\mu) U_\mu(x - a_\mu))] ]$$

Gauge transformation at  $x$  changes 2d link variables

$$U_\mu(x) \rightarrow \exp(-\alpha \epsilon \lambda_a G_a(x)) U_\mu(x)$$

$$U_\mu(x - a_\mu) \rightarrow U_\mu(x - a_\mu) \exp(\alpha \epsilon \lambda_a G_a(x))$$

Dynamical steps are interspersed with several gauge cooling steps



The strength of the cooling is determined by  
cooling steps  
gauge cooling parameter  $\alpha$

During cooling, unitarity norm decays to a minimum  
with a power law behaviour



# Adaptive cooling, Fourier accelerated cooling

[Aarts, Bongiovanni, Seiler, Sexty, Stamatescu (2013)]




Get to minimum quickest

Stepsize dependent on gradient  
Adaptive cooling

Low momentum modes cool slower  
Fourier accelerated cooling

See Lorenzo Bongiovanni's poster



# Polyakov chain model [Seiler, Sexty, Stamatescu (2012)]

exactly solvable toy model with gauge symmetry

$$S = -\beta_1 \text{Tr} U_1 \dots U_N - \beta_2 \text{Tr} U_N^{-1} \dots U_1^{-1} \quad U_i \in SU(3)$$

$$\beta_1 = \beta + \kappa e^\mu \quad \beta_2 = \beta^* + \kappa e^{-\mu}$$

Complex action for  $\kappa, \mu > 0$


Observables:  $\text{Tr} P^k$  with  $P = U_1 \dots U_N$

Averages independent of  $N$

Calculated with numerical integration at  $N=1$



Gauge symmetry

$$U_i \rightarrow V_i U_i V_{i+1}^{-1}$$



Smaller cooling  $\rightarrow$  excursions

“Skirt” develops  
small skirt gives correct result



# Heavy Quark QCD

Hopping parameter expansion of the fermion determinant  
Spatial hoppings are dropped

$$\text{Det } M(\mu) = \prod_x \text{Det} (1 + C P_x)^2 \text{Det} (1 + C' P_x^{-1})^2$$

$$P_x = \prod_\tau U_0(x + \tau a_0) \quad C = [2\kappa \exp(\mu)]^{N_\tau} \quad C' = [2\kappa \exp(-\mu)]^{N_\tau}$$

$$S = S_W[U_\mu] + \ln \text{Det } M(\mu)$$


De Pietri, Feo, Seiler, Stamatescu '07

Studied with reweighting

CLE study using gaugecooling

[Seiler, Sexty, Stamatescu (2012)]

See Nucu Stamatescu's poster



# Extension to full QCD with light quarks

[Sexty, arXiv:1307.7748]

QCD with staggered fermions

$$Z = \int DU e^{-S_G} \det M$$

$$M(x, y) = m \delta(x, y) + \sum_v \frac{\eta_v}{2a_v} (e^{\delta_{v4}\mu} U_v(x) \delta(x+a_v, y) - e^{-\delta_{v4}\mu} U_v^{-1}(x-a_v, y) \delta(x-a_v, y))$$

Still doubling present  $N_F=4$

$$Z = \int DU e^{-S_G} (\det M)^{N_F/4}$$

Langevin equation

$$U' = \exp(i\lambda_a (\epsilon i D_a S[U] + \sqrt{\epsilon} \eta_a)) U$$

$$K_{axv}^G = -D_{axv} S_G[U]$$

$$K_{axv}^F = \frac{N_F}{4} D_{axv} \ln \det M = \frac{N_F}{4} \text{Tr} (M^{-1} M'_{va}(x, y, z))$$


$$M'_{va}(x, y, z) = D_{azv} M(x, y)$$

Estimated using random sources  
1 CG solution per update

# Zero chemical potential


Drift is built from random numbers      real only on average


Cooling is essential already for small (or zero)  $\mu$



# Comparison of HQCD to full QCD

Qualitatively similar, chemical potential “rescaled”





## Conclusion


QCD = HQCD for quark mass  $> 4$

(For large mass) HQCD is qualitatively similar to QCD




# Average sign of the fermion determinant for small mass


Costly observable, only on small lattices possible



$$\langle \exp(2i\varphi) \rangle = \left\langle \frac{\det M(\mu)}{\det M(-\mu)} \right\rangle$$




# Horizontal slice of phase diagram



# Silver Blaze phenomenon

No dependence on chemical potential for small chemical potential

Zero temperature physics



Finite size effects important

Consistent with Silver Blaze

# Conclusions

New algorithm for Complex Langevin of gauge theories:  
Gauge cooling

Tested on exactly solvable toy model Polyakov chain  
Results for HQCD with heavy quarks with chemical potential  
Validated with reweighting

Results for full QCD with light quarks  
No sign or overlap problem  
CLE works all the way into saturation region  
Low temperatures are more demanding