

The Effect of the Low Energy Constants on the Spectral Properties of the Wilson Dirac Operator

Savvas Zafeiropoulos

July 29-August 3, 2013

Wilson Chiral Perturbation Theory

- Wilson term breaks χ - symmetry explicitly
- Lattice spacing effects lead to new terms in $\chi - PT$
Sharpe and Singleton (1998), Rupak and Shoresh (2002), Baer, Rupak and Shoresh (2004)
- ϵ - regime where in the thermodynamic, chiral and continuum limit $mV\Sigma$, $zV\Sigma$ and a^2VW_i kept fixed.
- At order a^2 it involves three Low Energy Constants (LECs)

$$Z_{N_f}(m, z; a) = \int_{\mathcal{M}} dU e^{-S[U]},$$

where the action is

$$S = -\frac{m}{2}\Sigma V \text{tr}(U + U^\dagger) - \frac{z}{2}\Sigma V \text{tr}(U - U^\dagger) \\ + a^2 V W_6 [\text{tr}(U + U^\dagger)]^2 + a^2 V W_7 [\text{tr}(U - U^\dagger)]^2 + a^2 V W_8 \text{tr}(U^2 + U^{\dagger 2}).$$

Wilson Chiral Perturbation Theory

- Wilson term breaks χ - symmetry explicitly
- Lattice spacing effects lead to new terms in $\chi - PT$
Sharpe and Singleton (1998), Rupak and Shoresh (2002), Baer, Rupak and Shoresh (2004)
- ϵ - regime where in the thermodynamic, chiral and continuum limit $mV\Sigma$, $zV\Sigma$ and a^2VW_i kept fixed.
- At order a^2 it involves three Low Energy Constants (LECs)

$$Z_{N_f}(m, z; a) = \int_{\mathcal{M}} dU e^{-S[U]},$$

where the action is

$$S = -\frac{m}{2}\Sigma V \text{tr}(U + U^\dagger) - \frac{z}{2}\Sigma V \text{tr}(U - U^\dagger) \\ + a^2 V W_6 [\text{tr}(U + U^\dagger)]^2 + a^2 V W_7 [\text{tr}(U - U^\dagger)]^2 + a^2 V W_8 \text{tr}(U^2 + U^{\dagger 2}).$$

Wilson Chiral Perturbation Theory

- Wilson term breaks χ - symmetry explicitly
- Lattice spacing effects lead to new terms in $\chi - PT$
Sharpe and Singleton (1998), Rupak and Shoresh (2002), Baer, Rupak and Shoresh (2004)
- ϵ - regime where in the thermodynamic, chiral and continuum limit $mV\Sigma$, $zV\Sigma$ and a^2VW_i kept fixed.
- At order a^2 it involves three Low Energy Constants (LECs)

$$Z_{N_f}(m, z; a) = \int_{\mathcal{M}} dU e^{-S[U]},$$

where the action is

$$S = -\frac{m}{2}\Sigma V \text{tr}(U + U^\dagger) - \frac{z}{2}\Sigma V \text{tr}(U - U^\dagger) \\ + a^2 VW_6 [\text{tr}(U + U^\dagger)]^2 + a^2 VW_7 [\text{tr}(U - U^\dagger)]^2 + a^2 VW_8 \text{tr}(U^2 + U^{\dagger 2}).$$

The Goal

- Facilitate simulations in the deep chiral regime by an exact, analytical understanding of the average behavior of the smallest eigenvalues
- Chiral symmetry breaking from lattice spacing
- Stability of lattice simulations

The Goal

- Facilitate simulations in the deep chiral regime by an exact, analytical understanding of the average behavior of the smallest eigenvalues
- Chiral symmetry breaking from lattice spacing
- Stability of lattice simulations

The Goal

- Facilitate simulations in the deep chiral regime by an exact, analytical understanding of the average behavior of the smallest eigenvalues
- Chiral symmetry breaking from lattice spacing
- Stability of lattice simulations

Introduction of the Model

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ Hermitian

- $B : (n + \nu) \times (n + \nu)$ Hermitian

- $W : n \times (n + \nu)$ Complex

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines

or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ Hermitian

- $B : (n + \nu) \times (n + \nu)$ Hermitian

- $W : n \times (n + \nu)$ Complex

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines

or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ **Hermitian**

- $B : (n + \nu) \times (n + \nu)$ **Hermitian**

- $W : n \times (n + \nu)$ **Complex**

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

$$\lambda_k^W \in \mathbb{R}$$

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ Hermitian

- $B : (n + \nu) \times (n + \nu)$ Hermitian

- $W : n \times (n + \nu)$ Complex

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines

or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ **Hermitian**

- $B : (n + \nu) \times (n + \nu)$ **Hermitian**

- $W : n \times (n + \nu)$ **Complex**

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

$$\lambda_k^W \in \mathbb{R}$$

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ **Hermitian**

- $B : (n + \nu) \times (n + \nu)$ **Hermitian**

- $W : n \times (n + \nu)$ **Complex**

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

$$\lambda_k^W \in \mathbb{R}$$

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ **Hermitian**

- $B : (n + \nu) \times (n + \nu)$ **Hermitian**

- $W : n \times (n + \nu)$ **Complex**

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

$$\lambda_k^W \in \mathbb{R}$$

Wilson Dirac operator and RMT

- Partition function of D_W with N_f flavors :

$$Z_{N_f}^{RMT,\nu} = \int dD_W \det^{N_f} (D_W + m) P(D_W)$$

- $P(D_W) \rightarrow$ is a Gaussian

- $D_W = \begin{pmatrix} aA & W \\ W^\dagger & aB \end{pmatrix} + am_6 + a\lambda_7\gamma_5$ (Damgaard et al (2010), Akemann et al (2010), Kieburg et al (2011,2012))

- $A : n \times n$ **Hermitian**

- $B : (n + \nu) \times (n + \nu)$ **Hermitian**

- $W : n \times (n + \nu)$ **Complex**

- m_6 and λ_7 scalar random variables

- At $a = 0$: D_W has ν generic zero modes

- At finite a : definition of the index through spectral flow lines

or equivalently $\nu = \sum_{\lambda_k^W \in \mathbb{R}} \text{sign}(\langle k | \gamma_5 | k \rangle)$ Itoh et al (1987)

- $D_W = \frac{1}{2}\gamma_\mu(\nabla_\mu + \nabla_\mu^*) - \frac{1}{2}a\nabla_\mu^*\nabla_\mu$
- At $a \neq 0$ is non-Hermitian but retains γ_5 -Hermiticity
 $D_W^\dagger = \gamma_5 D_W \gamma_5$
- Eigenvalues of D_W because of the γ_5 -Hermiticity occur in complex conjugate pairs or are real
- ONLY eigenvectors corresponding to real eigenvalues have non vanishing chirality $\langle k | \gamma_5 | k \rangle$


- $D_W = \frac{1}{2}\gamma_\mu(\nabla_\mu + \nabla_\mu^*) - \frac{1}{2}a\nabla_\mu^*\nabla_\mu$
- At $a \neq 0$ is non-Hermitian but retains γ_5 -Hermiticity
 $D_W^\dagger = \gamma_5 D_W \gamma_5$
- Eigenvalues of D_W because of the γ_5 -Hermiticity occur in complex conjugate pairs or are real
- ONLY eigenvectors corresponding to real eigenvalues have non vanishing chirality $\langle k | \gamma_5 | k \rangle$

- $D_W = \frac{1}{2}\gamma_\mu(\nabla_\mu + \nabla_\mu^*) - \frac{1}{2}a\nabla_\mu^*\nabla_\mu$
- At $a \neq 0$ is non-Hermitian but retains γ_5 -Hermiticity
 $D_W^\dagger = \gamma_5 D_W \gamma_5$
- Eigenvalues of D_W because of the γ_5 -Hermiticity occur in complex conjugate pairs or are real
- ONLY eigenvectors corresponding to real eigenvalues have non vanishing chirality $\langle k | \gamma_5 | k \rangle$

- $D_W = \frac{1}{2}\gamma_\mu(\nabla_\mu + \nabla_\mu^*) - \frac{1}{2}a\nabla_\mu^*\nabla_\mu$
- At $a \neq 0$ is non-Hermitian but retains γ_5 -Hermiticity
 $D_W^\dagger = \gamma_5 D_W \gamma_5$
- Eigenvalues of D_W because of the γ_5 -Hermiticity occur in complex conjugate pairs or are real
- ONLY eigenvectors corresponding to real eigenvalues have non vanishing chirality $\langle k | \gamma_5 | k \rangle$

The Eigenvalue Densities

Lattice results vs RMT




$$\hat{a}_6 = \hat{a}_7 = 0.25, \hat{a}_8 = 0.7$$

$$\hat{m} = 5.3$$

$\nu = 0$ (top) and

$\nu = 1$ (bottom)


(Deuzeman, Wenger and Wuilloud (2011))



$$\hat{m} = 4.8, \nu = 2$$


(Damgaard, Heller and Splittorff (2011))

Lattice results vs RMT



The density of real eigenvalues
of D_W

Damgaard, Heller and Splittorff (2012)



Cumulative eigenvalue
distributions of D_5 with all
 $W_{6/7/8}$ included at $\nu = 0$

(Deuzeman, Wenger and Wuilloud (2011))

The effects of W_6 and W_7 when $W_8 = 0$

- \hat{a}_6 and \hat{a}_7 introduced through the addition of the Gaussian stochastic variable $\hat{m}_6 + \hat{\lambda}_7 \gamma_5$ to D_W
- $D = D_W + (m + \hat{m}_6)\mathbf{1} + \hat{\lambda}_7 \gamma_5$
- When $\hat{a}_8 = 0$ D_W is anti-Hermitian,
- the eigenvalues of $D_W(\hat{\lambda}_7, \hat{m}_6) = D - m$ are given by

$$\hat{z}_{\pm} = \hat{m}_6 \pm i\sqrt{\lambda_W^2 - \hat{\lambda}_7^2}$$

where $i\lambda_W$ is an eigenvalue of D_W

The effects of W_6 and W_7 when $W_8 = 0$

- \hat{a}_6 and \hat{a}_7 introduced through the addition of the Gaussian stochastic variable $\hat{m}_6 + \hat{\lambda}_7 \gamma_5$ to D_W
- $D = D_W + (m + \hat{m}_6)\mathbf{1} + \hat{\lambda}_7 \gamma_5$
- When $\hat{a}_8 = 0$ D_W is anti-Hermitian,
- the eigenvalues of $D_W(\hat{\lambda}_7, \hat{m}_6) = D - m$ are given by

$$\hat{z}_{\pm} = \hat{m}_6 \pm i\sqrt{\lambda_W^2 - \hat{\lambda}_7^2}$$

where $i\lambda_W$ is an eigenvalue of D_W

The effects of W_6 and W_7 when $W_8 = 0$

- \hat{a}_6 and \hat{a}_7 introduced through the addition of the Gaussian stochastic variable $\hat{m}_6 + \hat{\lambda}_7 \gamma_5$ to D_W
- $D = D_W + (m + \hat{m}_6)\mathbf{1} + \hat{\lambda}_7 \gamma_5$
- When $\hat{a}_8 = 0$ D_W is anti-Hermitian,
- the eigenvalues of $D_W(\hat{\lambda}_7, \hat{m}_6) = D - m$ are given by

$$\hat{z}_{\pm} = \hat{m}_6 \pm i\sqrt{\lambda_W^2 - \hat{\lambda}_7^2}$$

where $i\lambda_W$ is an eigenvalue of D_W


The effects of W_6 and W_7 when $W_8 = 0$

- \hat{a}_6 and \hat{a}_7 introduced through the addition of the Gaussian stochastic variable $\hat{m}_6 + \hat{\lambda}_7 \gamma_5$ to D_W
- $D = D_W + (m + \hat{m}_6)\mathbf{1} + \hat{\lambda}_7 \gamma_5$
- When $\hat{a}_8 = 0$ D_W is anti-Hermitian,
- the eigenvalues of $D_W(\hat{\lambda}_7, \hat{m}_6) = D - m$ are given by

$$\hat{z}_{\pm} = \hat{m}_6 \pm i\sqrt{\lambda_W^2 - \hat{\lambda}_7^2}$$


where $i\lambda_W$ is an eigenvalue of D_W

The effects of W_6 and W_7




Schematic plots of the effects of W_6 (left plot) and of W_7 (right plot). W_6 broadens the spectrum parallel to the real axis according to a Gaussian with width $4\hat{a}_6$, but does not change the continuum spectrum in a significant way. When $W_7 \neq 0$ and $W_6 = 0$ the purely imaginary eigenvalues invade the real axis through the origin and only the real (green crosses) are broadened by a Gaussian with width $4\hat{a}_7$

Distribution of additional real modes for $\nu = 1$




Notice that the two curves for $\hat{a}_7 = \hat{a}_8 = 0.1$ (right plot) are two orders smaller than the other curves (left plot). Notice the soft repulsion of the additional real modes from the origin at large \hat{a}_7 . The parameter \hat{a}_6 smooths the distribution.

Log-Log plots of additional real modes vs \hat{a} for $\nu = 0, 2$




Log-log plots of N_{add} as a function of \hat{a}_8 for $\nu = 0$ (left plot) and $\nu = 2$ (right plot). W_6 has no effect on N_{add} . Saturation around zero due to a non-zero value of \hat{a}_7 . For $\hat{a}_7 = 0$ (lowest curves) the average number of additional real modes behaves like $\hat{a}_8^{2\nu+2}$. Kieburg, Verbaarschot and SZ (2011)

Distribution of additional real modes for $\hat{a} \gg 1$




At $\hat{a} \gg 1$ ρ_r develops square root singularities at the boundaries. Finite matrix size + finite lattice spacing $\rightarrow \rho_r$ has a tail dropping off much faster than the size of the support. The dependence on W_6 and ν is completely lost.

Projected distribution of the complex eigenvalues for $\nu = 1$



The distribution of the complex eigenvalues projected onto the imaginary axis for $\nu = 1$. Notice that \hat{a}_6 does not affect this distribution. The comparison of $\hat{a}_7 = \hat{a}_8 = 0.1$ with the continuum result (black curve) shows that ρ_{cp} is still a good quantity to extract the chiral condensate Σ at small lattice spacing.

Chirality distribution for $\nu = 1$



The distribution is symmetric around the origin. At small \hat{a}_8 the distributions for $(\hat{a}_6, \hat{a}_7) = (1, 0.1), (0.1, 1)$ are almost the same Gaussian as the analytical result predicts. At large \hat{a}_8 the maximum reflects the predicted square root singularity which starts to build up. We have not included the case $\hat{a}_{6/7/8} = 0.1$ since it exceeds the other curves by a factor of 10 to 100.

Extracting the LECs of Wilson chPT

Extracting the LECs of Wilson chPT

$$\begin{aligned}
 & \left. K_1^{(n)} \left(\frac{\bar{z}_1}{2n}, \frac{\bar{z}_2}{2n} \right) \right|_{\bar{z}_1 \leq 1} \sim \frac{n^2}{4(2n)^{3/2}} f(\varphi_1, \varphi_2) \dots \times \prod_{j=1}^n |K_1^{(n)}(z_j^{(1)}, z_j^{(2)})| K_2^{(n)}(z_j^{(1)}, z_j^{(2)}) |K_3^{(n)}(m_j, z_j^{(1)})| \dots \\
 & g_2(z_1, z_2) = g_1(z_1, z_2) \delta(y_1) \delta(y_2) + g_2(z_1) \delta(x_1 - x_2) \delta(y_1 + y_2), \quad (17) \quad \frac{1}{(m_1 + 2n + \nu)^2} \int_{\mathbb{R}^2} \prod_{l=1}^{N_0} \det(Z + m_l + \nu \sigma_l) \\
 & \times \exp \left[\sum_{j=1}^2 \left(-4g_0^2 \cos^2 \frac{k(\bar{z}_j, \alpha)}{\sqrt{\bar{z}_j, \alpha}} \right) \times \exp \left[-\frac{n}{4a^2} (x_1 + x_2 - \frac{a^2(\mu_1 + \mu_2)}{n})^2 + \frac{n}{4} (x_1 - x_2)^2 \right] \right] \quad (18) \\
 & \times \left[\operatorname{sign}(x_1 - x_2) - \operatorname{erf} \left[\sqrt{\frac{n(1+a^2)}{4a^2}} (x_1 - x_2) - \sqrt{\frac{a^2}{4n(1+a^2)}} (\mu_1 - \mu_2) \right] \right] \cdot \frac{d\alpha_j^{(1)} d\alpha_j^{(2)}}{K_2^{(n)}(z_j^{(1)}, z_j^{(2)})} \prod_{j=k_1+1}^{n-k_2} d\alpha_j^{(1)} d\alpha_j^{(2)}. \quad (C.1) \\
 & g_1(z) = -2 \operatorname{sign}(y) \exp \left[-\frac{n}{a^2} \left(x - \frac{a^2(\mu_1 + \mu_2)}{2n} \right)^2 - ny^2 \right], \quad (19) \\
 & g_3(x) = \exp \left[-\frac{n}{2a^2} \left(x - \frac{a^2 \mu_1}{n} \right)^2 \right], \quad (20) \\
 & \times \left[\operatorname{sign}(x_2) - \operatorname{erf} \left[\frac{-x_2}{\sqrt{8a^2}} \right] \right] \\
 & + 8na^2 \operatorname{sign} \alpha_1 \exp \left[-\frac{1}{8a^2} (\bar{r}_1 - \bar{w}_a)^2 \right] \Delta K_1^{(n)} \left(\frac{\bar{z}_1}{2n}, \frac{\bar{z}_1}{2n} \right). \quad \left. \frac{-x_1}{-k_2 \bar{z} = 0} \right]
 \end{aligned}$$

- the average number of the additional real modes for the lowest index:

$$N_{\text{add}}^{\nu=0} \stackrel{\bar{a} \leq 1}{\approx} 2V \bar{a}^2 (W_8 - 2W_7), \quad (75)$$

- the width of the Gaussian shaped strip of complex eigenvalues:

$$2\sigma \stackrel{\bar{a} \leq 1}{\approx} 4\bar{a} \sqrt{\frac{W_6 - 2W_6}{V\Sigma^2}}, \quad (76)$$

- the variance of the distribution of chirality over the real eigenvalues:

$$\langle (V\Sigma \bar{x})^2 \rangle_{\rho_x} \stackrel{\bar{a} \leq 1}{\approx} 8V \bar{a}^2 (\nu W_6 - W_6 - W_7), \quad \nu > 0. \quad (77)$$



(Figure courtesy of M. Kieburg)

Conclusions

- Studied the effect of the three LECs on the spectrum of D_W .
- W_6 and W_7 can be interpreted as collective fluctuations of the spectrum while W_8 induces interactions among all modes.
- Analytical and numerical results of the eigenvalue densities of D_W
- At small lattice spacing we propose the following quantities for the extraction of LECs

$$\tilde{a}^2 V \begin{bmatrix} 0 & -2 & 1 \\ -2 & 0 & 1 \\ -1 & -1 & 1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} W_6 \\ W_7 \\ W_8 \end{bmatrix} = \frac{\pi^2}{8} \begin{bmatrix} 4N_{\text{add}}^{\nu=0} / \pi^2 \\ 2\sigma^2 / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=1} / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=2} / \Delta^2 \end{bmatrix}$$

Conclusions

- Studied the effect of the three LECs on the spectrum of D_W .
- W_6 and W_7 can be interpreted as collective fluctuations of the spectrum while W_8 induces interactions among all modes.
- Analytical and numerical results of the eigenvalue densities of D_W
- At small lattice spacing we propose the following quantities for the extraction of LECs

$$\tilde{a}^2 V \begin{bmatrix} 0 & -2 & 1 \\ -2 & 0 & 1 \\ -1 & -1 & 1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} W_6 \\ W_7 \\ W_8 \end{bmatrix} = \frac{\pi^2}{8} \begin{bmatrix} 4N_{\text{add}}^{\nu=0} / \pi^2 \\ 2\sigma^2 / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=1} / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=2} / \Delta^2 \end{bmatrix}$$

Conclusions

- Studied the effect of the three LECs on the spectrum of D_W .
- W_6 and W_7 can be interpreted as collective fluctuations of the spectrum while W_8 induces interactions among all modes.
- Analytical and numerical results of the eigenvalue densities of D_W
- At small lattice spacing we propose the following quantities for the extraction of LECs

$$\tilde{a}^2 V \begin{bmatrix} 0 & -2 & 1 \\ -2 & 0 & 1 \\ -1 & -1 & 1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} W_6 \\ W_7 \\ W_8 \end{bmatrix} = \frac{\pi^2}{8} \begin{bmatrix} 4N_{\text{add}}^{\nu=0} / \pi^2 \\ 2\sigma^2 / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=1} / \Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_X}^{\nu=2} / \Delta^2 \end{bmatrix}$$

Conclusions

- Studied the effect of the three LECs on the spectrum of D_W .
- W_6 and W_7 can be interpreted as collective fluctuations of the spectrum while W_8 induces interactions among all modes.
- Analytical and numerical results of the eigenvalue densities of D_W
- At small lattice spacing we propose the following quantities for the extraction of LECs

$$\tilde{a}^2 V \begin{bmatrix} 0 & -2 & 1 \\ -2 & 0 & 1 \\ -1 & -1 & 1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} W_6 \\ W_7 \\ W_8 \end{bmatrix} = \frac{\pi^2}{8} \begin{bmatrix} 4N_{\text{add}}^{\nu=0}/\pi^2 \\ 2\sigma^2/\Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_\chi}^{\nu=1}/\Delta^2 \\ \langle \tilde{x}^2 \rangle_{\rho_\chi}^{\nu=2}/\Delta^2 \end{bmatrix}$$

Stay Tuned!



for upcoming results ...

Thank you for your attention!

Collaborators:

Mario Kieburg 8E 17.30

A classification of 2– dim Lattice Theory

Jacobus Verbaarschot 7D 15.40

Discretization Effects in the ϵ Domain of QCD